



# The Effect of Polystyrene Microplastic Exposure in the Rearing Water on Muscle Morphology of Mutiara Catfish (*Clarias gariepinus* Burchell, 1822)

Nurul Suwartiningsih<sup>1</sup>, Aulia Fayzha Aszhara<sup>1</sup>, Rani Marsela<sup>1</sup>, Dian Eka Wijayanti<sup>2</sup>

<sup>1</sup> Study Program of Biology, Faculty of Applied Science and Technology, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

<sup>2</sup> Study Program of Mathematics, Faculty of Applied Science and Technology, Universitas Ahmad Dahlan, Yogyakarta, Indonesia.

Received: May 30, 2025

Revised: July 21, 2025

Accepted: July 27, 2025

Published: July 31, 2025

Corresponding Author:

Nurul Suwartiningsih

[nurul.suwartiningsih@bio.uad.ac.id](mailto:nurul.suwartiningsih@bio.uad.ac.id)

DOI: [10.29303/jppipa.v11i7.11537](https://doi.org/10.29303/jppipa.v11i7.11537)

© 2025 The Authors. This open access article is distributed under a (CC-BY License)



**Abstract:** Polystyrene (PS) is one of the most abundant microplastics in freshwater and can accumulate in fish tissues. This study investigated the effect of PS microplastic exposure in rearing water on the abundance of microplastics in muscle, and the diameter and area of myofibers in Mutiara catfish (*Clarias gariepinus*). A total of 120 fish were exposed to PS microplastics at concentrations of 0, 1, 10, and 100 mg/L for 28 days. Muscle samples from the right abdomen were processed using the paraffin method and stained with hematoxylin-eosin. Microplastics were extracted using 10% KOH and observed under a stereo microscope. Myofiber diameter and area were measured with Image Raster software. Data were analyzed using the Kruskal-Wallis test. Results showed significant differences among treatments ( $P<0.05$ ) for all parameters. The control group had the lowest microplastic abundance and the largest myofiber size, while the highest accumulation occurred at 100 mg/L, and the smallest myofiber diameter and area were found at 10 mg/L. Exposure to PS microplastics leads to accumulation in muscle tissue and a reduction in myofiber dimensions. These findings highlight the potential adverse impacts of microplastics on fish health and underline the importance of further studies regarding ecological and human health implications.

**Keywords:** Microplastic exposure; Muscle tissue; Myofiber area; Myofiber diameter; Polystyrene toxicity

## Introduction

Microplastics are increasingly recognized as emerging pollutants in aquatic ecosystems, appearing in various polymers, shapes, and concentrations (Smith et al., 2018). These particles originate primarily from consumer products such as plastic bags, beverage bottles, personal care items, and cleaning agents (Nainggolan et al., 2022). Due to their diverse polymer composition, ranging from high to low density plastics, microplastics are easily transported by water currents and accumulate in both water columns and sediments (Owowenu et al., 2023). Numerous studies have

reported the detection of microplastics in freshwater ecosystems, particularly in river areas (Dhea et al., 2023; Lofty et al., 2023). Research has identified microplastics in the Pekalongan River estuary, with abundances ranging from 45.2 to 99.1 particles L<sup>-1</sup> in surface water, dominated by fragment and film morphologies (Ismanto et al., 2023). Microplastics pose ecological risks by negatively impacting aquatic organisms and degrading environmental quality (Li et al., 2023; Yuranda et al., 2024).

Rivers contaminated with microplastics may transfer these particles into aquaculture systems (Santikanuri et al., 2025), thereby affecting farmed fish.

### How to Cite:

Suwartiningsih, N., Aszhara, A. F., Marsela, R., & Wijayanti, D. E. The Effect of Polystyrene Microplastic Exposure in the Rearing Water on Muscle Morphology of Mutiara Catfish (*Clarias gariepinus* Burchell, 1822). *Jurnal Penelitian Pendidikan IPA*, 11(7), 295-302. <https://doi.org/10.29303/jppipa.v11i7.11537>

Microplastic contamination in aquaculture environments can deteriorate water quality, compromise fish health, and ultimately reduce aquaculture productivity (Filho et al., 2023). Research findings indicate that tilapia (*Oreochromis niloticus*) reared in the Pelayaran River, Sidoarjo, contain microplastic residues, attributed to pollution in the surrounding aquatic environment (Al-Fatih, 2022).

Among the various types of microplastics, polystyrene is commonly detected in aquatic habitats (Jones et al., 2020) originating from food and beverage packaging as well as the automotive industry (Nurhadi et al., 2017). Research on the Mahakam River found that polystyrene accounted for 6.45% of the total plastic waste, with microplastics comprising 12.5% of the total abundance (Kurniawan et al., 2023). Environmental exposure, especially ultraviolet radiation and weathering, accelerates the breakdown of polystyrene into microplastic fragments (Hwang et al., 2020).

Microplastics can be ingested by fish through their gills, digestive tract and muscles (Sawalman et al., 2021). Microplastics present in water can enter fish muscles via the bloodstream (Akhbarizadeh et al., 2018). When microplastics enter the digestive tract, they can be transported through the blood vessels and eventually accumulate in muscle tissue (Liu et al., 2022). Previous research has shown that exposure to microplastics at a concentration of 100 mg/L in tilapia (*Oreochromis niloticus*) can cause muscle damage (Hamed et al., 2021). One of the commonly cultivated and consumed fish species is Mutiara catfish (*Clarias gariepinus* Burchell, 1822) (Iswanto et al., 2016). Despite growing concern over microplastic contamination, the histopathological effects of polystyrene microplastics on muscle morphology in Mutiara catfish remain understudied. Most research has focused on gastrointestinal tissues, with limited attention to muscle damage in economically important aquaculture species under controlled exposure.

The novelty of this research lies in evaluating the sub-lethal and tissue-specific effects of polystyrene microplastic exposure on Mutiara catfish muscle morphology. This study provides critical insight into how microplastics may compromise fish meat quality, a concern for both environmental health and food safety. Therefore, this research is important to understand potential histological damage caused by microplastic contamination in aquaculture systems. The findings may inform sustainable fish farming practices, guide risk assessments of microplastic exposure in aquatic food chains, and support the development of environmental policies to mitigate plastic pollution impacts on food-producing aquatic ecosystems.

## Method

### Research Time and Location

This research was conducted from January to November 2024. Maintenance and treatment took place in Dengok Kulon, Bugisan, Prambanan, Klaten, while data preparation, collection, and analysis were carried out at the Laboratory of Animal Structure and Physiology, Universitas Ahmad Dahlan.

### Experimental Design

This study employed an experimental approach using a completely randomized design (CRD) with four treatment groups: a control group (0 mg/L polystyrene microplastic), T1 (1 mg/L), T2 (10 mg/L), and T3 (100 mg/L). Each treatment was replicated six times using separate rearing tanks, with five Mutiara catfish (*Clarias gariepinus*) allocated per tank, resulting in a total of 120 fish. The exposure period lasted for 28 days. The parameters observed in this study included the abundance of microplastics in muscle tissue, the diameter of myofibers, and the area of myofibers.

### Microplastics Exposure

Microplastic preparation was carried out by pulverizing polystyrene plastic with a blender, followed by filtering through a 35-mesh sieve. The microplastics were weighed according to the treatment concentrations. Mutira catfish were acclimatized for ten days by being given food at 08.00 and 16.00 WIB, equivalent to 3% of their body weight (Tang et al., 2024). Microplastic exposure was conducted during water changes every two days for 28 days, based on the designated exposure concentrations.

### Muscle Tissue Preparation

At the end of the treatment, the fish were fasted for 24 hours before being weighed and their organs harvested on day 29. Muscle tissues were weighed and fixed in a 10% formalin solution (Zulfadhl et al., 2016). The preparation process included dehydration with graded alcohol (70% to absolute), de-alcoholization with toluol solution infiltration, embedding in paraffin, sectioning with a microtome, fixation using Mayer's albumin, and staining with hematoxylin-eosin. The prepared tissue slides were examined under a light microscope at 40x and 100x magnification, with observations made in five fields of view per slide. Documentation was carried out using a Beta View camera system. Muscle parameters, including myofiber diameter and area, were measured using Image Raster.

### Microplastic Extraction

Microplastic abundance analysis was carried out starting with muscle organs placed in 30 mL flask bottles

containing a 10% potassium hydroxide (KOH) solution. The samples were left at room temperature for three days to allow digestion. After three days, the contents of the flask bottles were poured into empty bottles equipped with a funnel and filter paper. The filter paper was left to dry completely before distilled water was added to dissolve any microplastics adhering to the filter. The resulting solution was examined under a stereo microscope at 4x magnification, and the microplastic abundance was quantified (Suwartiningsih et al., 2023).

#### Data Analysis

The raw data were initially processed using Microsoft Excel, followed by statistical analysis with IBM SPSS version 20. Differences between treatments were considered significant at  $P < 0.05$  for all measured parameters and were evaluated using a one-way analysis of variance (ANOVA). When significant differences were detected, further analysis was conducted using Duncan's Multiple Range Test (DMRT).

## Result and Discussion

#### Microplastic Abundance

The results showed that the abundance of microplastics in the muscles of Mutiara catfish was lowest in the control group ( $0.00 \pm 0.00$  particles/g) and highest in the T3 ( $2.63 \pm 2.72$  particles/g). Statistical analysis indicated that T3 was significantly different from the control and other treatment groups ( $P < 0.05$ ). Regression analysis demonstrated that exposure to polystyrene microplastics in rearing water for 28 days significantly affected the abundance of microplastics in the muscles of Mutiara catfish ( $P < 0.05$ ). This suggests a positive correlation, where higher concentrations of microplastics in the rearing water led to greater microplastic accumulation in the fish muscles, as shown in Table 1.

The research showed that microplastic exposure starting from 1 mg/L in the rearing media, the abundance of microplastics in the Mutiara catfish muscle was also high. Previous research found that exposure to nano plastic polystyrene (NPPs) at a concentration of 1 mg/L for three days in zebrafish (*Danio rerio*) resulted in NPPs accumulation of  $49 \pm 17$  mg/g of muscle tissue (Chen et al., 2017). The high abundance of microplastics in fish organs is influenced by the increasing concentration of microplastic particles in the water (Popi et al., 2025; Rofiq & Sari, 2022; Wahid & Joesidawati, 2024), and vice versa (Fitriyani et al.,

2025) allowing microplastics to enter the fish through food ingestion (Roch et al., 2020; Yona et al., 2022).

**Table 1.** Polystyrene microplastic abundance in Mutiara catfish muscle


Treatment	Microplastic abundance (particles/g)
Control	$0.00 \pm 0.00^a$
T1	$0.85 \pm 1.48^{ab}$
T2	$0.74 \pm 1.28^{ab}$
T3	$2.63 \pm 2.72^b$

Description: superscript different letters indicate there is a significant difference ( $P < 0.05$ ) and vice versa

The detection of microplastics in the muscles of Mutiara catfish in this study indicates bioaccumulation, or the accumulation of contaminant particles in the body (Miller et al., 2020; Utomo & Muzaki, 2022). When microplastics enter the gastrointestinal tract, larger particles may settle or be excreted (Purnama et al., 2021) in feces. However, as microplastics break down into smaller particles, they can travel through the bloodstream (Nurwahyunani et al., 2022) and accumulate in muscle tissue (Yona et al., 2021). Particles around 5  $\mu\text{m}$  in size can infiltrate the gills, intestines, and liver, while microplastics approximately 20  $\mu\text{m}$  in size can penetrate the gills and intestines (Guanting et al., 2021). Particles smaller than 100  $\mu\text{m}$  are capable of reaching muscle tissue (Barboza et al., 2020).

#### Diameter and Area of Mutiara Catfish Myofiber

The results showed that the diameter of myofiber in Mutiara catfish muscle was lowest in T2 ( $83.58 \pm 25.01$   $\mu\text{m}$ ) and highest in control ( $122.11 \pm 40.06$   $\mu\text{m}$ ). The area of myofiber in Mutiara catfish muscle was lowest in T2 ( $5.96 \pm 3.77 \times 10^3$   $\mu\text{m}^2$ ) and highest in control ( $12.94 \pm 8.65 \times 10^3$   $\mu\text{m}^2$ ) (Table 2). Statistical analysis revealed that both myofiber diameter and area were significantly different between control and the other treatment groups ( $P < 0.05$ ). However, there were no significant differences in myofiber diameter and area among the P1, P2, and P3 groups. Regression analysis indicated that polystyrene microplastic exposure in the rearing water for 28 days significantly affected both the myofiber diameter and area in Mutiara catfish ( $P < 0.05$ ). These findings are consistent with existing theories which suggest that exposure to toxic substances such as polystyrene microplastics can impair myogenesis and induce muscle atrophy, leading to a significant reduction in myofiber diameter and area.



**Figure 1.** Diameter and area of Mutiara catfish myofiber after 28 days exposure to polystyrene microplastics: (a) control (0 mg/L); (b) T1 (1 mg/L); (c) T2 (10 mg/L); and (d) T3(100 mg/L). Blue arrows indicate normal myofiber diameter and red arrows indicate decreased myofiber diameter (hematoxylin-eosin staining; scale bar 100  $\mu\text{m}$ )

**Table 2.** Diameter and area of Mutiara catfish myofiber

Treatment	Diameter of myofiber ( $\mu\text{m}$ )	Area of myofiber ( $\times 10^3 \mu\text{m}^2$ )
Control	122.11 $\pm$ 40.06 <sup>b</sup>	12.49 $\pm$ 8.65 <sup>b</sup>
T1	87.80 $\pm$ 26.81 <sup>a</sup>	6.60 $\pm$ 4.87 <sup>a</sup>
T2	83.58 $\pm$ 25.01 <sup>a</sup>	5.96 $\pm$ 3.77 <sup>a</sup>
T3	86.32 $\pm$ 18.21 <sup>a</sup>	6.10 $\pm$ 2.52 <sup>a</sup>

Description: superscript different letters indicate there is a significant difference ( $P < 0.05$ ) and vice versa

The results in Table 2 indicate that Mutiara catfish exposed to polystyrene microplastics for 28 days experienced a reduction in both myofiber diameter and area. The regulation of myofiber diameter is influenced by several factors including myogenesis, muscle atrophy and apoptosis (Zhang et al., 2023). Previous research demonstrated that exposure to polystyrene microplastics (PS-MPs) at a concentration of 1000  $\mu\text{g}/\text{L}$  for 7 days in goldfish larvae (*Carassius auratus*) caused damage to mesenchymal muscle cells and led to muscle fiber atrophy (Yang et al., 2020). The changes in myofiber diameter can be observed in Figure 1.

Changes in muscle tissue can occur due to the introduction of contaminants in the muscle (Di Giacinto et al., 2023; Setyono et al., 2024). Styrene is a benzene-derived compound that has high toxicity (Aldi, 2024; Majid et al., 2018). Previous research revealed that exposure to contaminants at concentrations of 44  $\mu\text{g}/\text{kg}$

and 234  $\mu\text{g}/\text{kg}$  in Yellow catfish (*Pelteobagrus fulvidraco*) over eight weeks led to a reduction in muscle myofiber diameter (Zhang et al., 2021). Another study showed that exposure of fonofos contaminants at a concentration of 2 mg/L for 96 hours caused a decrease in myofiber diameter in Zebrafish (*Danio rerio*) (Arman & Üçüncü, 2021).

This study found a decrease in both myofiber diameter and myofiber area, thus disrupting the growth of muscle myofiber (Moenek & Toelle, 2021). Myofiber growth that is not optimal will have an impact on the elasticity and compactness of the meat (Suwiti et al., 2015; Zhang et al., 2021). Moreover, muscle injuries may serve as an indicator of exposure to environmental contaminants (Ciamarro et al., 2015). Among such contaminants, microplastics pose a significant toxic risk to all organisms (Aulia et al., 2023; Stapleton & Hai, 2023).

## Conclusion

This study demonstrated that exposure to polystyrene (PS) microplastics in rearing water for 28 days significantly increased microplastic accumulation in muscle tissue and reduced both myofiber diameter and area in Mutiara catfish (*Clarias gariepinus*). The highest accumulation was observed at a concentration of

100 mg/L, reaching 2.63 particles/g, indicating that higher concentrations of microplastics in water lead to greater accumulation in muscle. This trend reflects patterns observed in other aquatic species, where microplastic presence in the environment directly influenced the level of contamination in fish organs. Histological analysis also showed that fish in the treatment groups experienced significant reductions in myofiber diameter and area compared to the control. The control group exhibited average myofiber diameters of 122.11  $\mu\text{m}$  and areas of  $12.49 \times 10^3 \mu\text{m}^2$ , while exposed groups showed reduced diameters as low as 83.58  $\mu\text{m}$  and areas down to  $5.96 \times 10^3 \mu\text{m}^2$ . Similar muscle degradation patterns have been documented in other fish exposed to plastic contaminants and toxic substances, such as atrophied muscle fibers, disorganized tissue structures, and inhibited muscle growth, all of which impact meat quality. These results confirm that even short-term exposure to PS microplastics at environmentally relevant concentrations can lead to bioaccumulation and muscle deterioration. The findings emphasize the potential ecological risks of microplastics and their implications for food safety, particularly in aquaculture systems. This highlights the urgent need for improved environmental monitoring and regulatory policies to reduce microplastic pollution and safeguard both aquatic ecosystems and public health.

#### Acknowledgments

Acknowledgments are extended to Lembaga Penelitian dan Pengabdian kepada Masyarakat Universitas Ahmad Dahlan (LPPM UAD) for providing funding for the implementation of this research.

#### Author Contributions

Conceptualization, N.S. and A.F.A.; methodology, N.S. and A.F.A.; validation, N.S.; formal analysis, A.F.A.; investigation, A.F.A. and D.E.W.; resources, N.S.; data curation, A.F.A.; writing—original draft preparation, A.F.A.; writing—review and editing, N.S., A.F.A., and D.E.W.; visualization, A.F.A.; supervision, N.S. All authors have read and approved the final version of the manuscript.

#### Funding

This research was funded by Lembaga Penelitian dan Pengabdian kepada Masyarakat Universitas Ahmad Dahlan (LPPM UAD), grant number PD-045/SP3/LPPM-UAD/XI/2024 dated November 28, 2024.

#### Conflicts of Interest

We hereby affirm that the preparation of this article was conducted without any conflict of interest that could compromise the objectivity or integrity of the findings.

## References

Akhbarizadeh, R., Moore, F., & Keshavarzi, B. (2018). Investigating a probable relationship between microplastics and potentially toxic elements in fish muscles from northeast of Persian Gulf. *Environmental Pollution*, 232, 154–163. <https://doi.org/10.1016/j.envpol.2017.09.028>

Al-Fatih, A. N. F. (2022). Identifikasi mikroplastik pada sistem pencernaan ikan Nila (*Oreochromis niloticus*) di kali pelayaran Kabupaten Sidoarjo. *Environmental Pollution Journal*, 1(3), 237–244. <https://doi.org/10.58954/epj.v1i3.63>

Aldi, D. (2024). Dampak pencemaran benzena terhadap lingkungan serta mekanismenya dalam memicu kanker: Sebuah tinjauan literatur. *Jurnal Pengelolaan Lingkungan Berkelanjutan (Journal of Environmental Sustainability Management)*, 8(1), 87–98. <https://doi.org/10.36813/jplb.8.1.87-98>

Arman, S., & Üçüncü, S. I. (2021). Impact of acute fonofos exposure on skeletal muscle of Zebrafish: Histopathological and biometric analyses. *Annales de Limnologie*, 57(25). <https://doi.org/10.1051/limn/2021023>

Aulia, A., Azizah, R., Sulistyorini, L., & Rizaldi, M. A. (2023). Literature Review: Dampak mikroplastik terhadap lingkungan pesisir, biota laut dan potensi risiko kesehatan. *Jurnal Kesehatan Lingkungan Indonesia*, 22(3), 328–341. <https://doi.org/10.14710/jkli.22.3.328-341>

Barboza, L. G. A., Lopes, C., Oliveira, P., Bessa, F., Otero, V., Henriques, B., Raimundo, J., Caetano, M., Vale, C., & Guilhermino, L. (2020). Microplastics in wild fish from North East Atlantic Ocean and its potential for causing neurotoxic effects, lipid oxidative damage, and human health risks associated with ingestion exposure. *Science of the Total Environment*, 717, 134625. <https://doi.org/10.1016/j.scitotenv.2019.134625>

Chen, Q., Yin, D., Jia, Y., Schiwy, S., Legradi, J., Yang, S., & Hollert, H. (2017). Enhanced uptake of BPA in the presence of nanoplastics can lead to neurotoxic effects in adult Zebrafish. *Science of the Total Environment*, 609, 1312–1321. <https://doi.org/10.1016/j.scitotenv.2017.07.144>

Ciamarro, C. M., Pereira, B. F., Alves, R. M. da S., Valim, J. R. T., Pitol, D. L., & Caetano, F. H. (2015). Changes in muscle and collagen fibers of fish after exposure to urban pollutants and biodegradable detergents. *Microscopy Research*, 03(03), 33–40. <https://doi.org/10.4236/mr.2015.33005>

Dhea, L. A., Kurniawan, A., Ulfa, S. M., & Karimah, K. (2023). Correlation of microplastic size distribution and water quality parameters in the upstream

Brantas river. *Jurnal Penelitian Pendidikan IPA*, 9(2), 520-526.  
<https://doi.org/10.29303/jppipa.v9i2.2777>

Di Giacinto, F., Di Renzo, L., Mascilongo, G., Notarstefano, V., Gioacchini, G., Giorgini, E., Bogdanović, T., Petričević, S., Listić, E., Brkljača, M., Conti, F., Profico, C., Zambuchini, B., Di Francesco, G., Giansante, C., Diletti, G., Ferri, N., & Berti, M. (2023). Detection of microplastics, polymers and additives in edible muscle of Swordfish (*Xiphias gladius*) and Bluefin tuna (*Thunnus thynnus*) caught in the Mediterranean Sea. *Journal of Sea Research*, 192(January). <https://doi.org/10.1016/j.seares.2023.102359>

Filho, J. V. D., Pedroti, V. P., Santos, B. L. T., Pinheiro, M. M. de L., de Mira, Á. B., da Silva, F. C., e Silva, E. C. S., Cavali, J., Guedes, Elica Amara Cecilia Schons, S., & Vargas, D. (2023). First evidence of microplastics in freshwater from fish farms in Rondônia state, Brazil. *Helijon*, 9(4), 1-11. <https://doi.org/10.1016/j.helijon.2023.e15066>

Fitriyani, F., Eryati, R., & Ritonga, I. R. (2025). Analisis kelimpahan mikroplastik di dalam saluran pencernaan ikan Bawis (*Siganus canaliculatus*) hasil tangkapan nelayan lokal di perairan kota Bontang. *Jurnal Ilmu Lingkungan*, 23(3), 784-791. <https://doi.org/10.14710/jil.23.3.784-791>

Guanting, L., Ito, T., Kijima, Y., Yoshitake, K., Asakawa, S., Watabe, S., & Kinoshita, S. (2021). Zebrafish *Danio rerio* trunk muscle structure and growth from a spatial transcriptomics perspective. *BioRxiv*, 6. <https://doi.org/10.1101/2021.06.14.447777>

Hamed, M., Soliman, H. A. M., Badrey, A. E. A., & Osman, A. G. M. (2021). Microplastics induced histopathological lesions in some tissues of Tilapia (*Oreochromis niloticus*) early juveniles. *Tissue and Cell*, 71(February), 101512. <https://doi.org/10.1016/j.tice.2021.101512>

Hwang, J., Choi, D., Han, S., Jung, S. Y., Choi, J., & Hong, J. (2020). Potential toxicity of polystyrene microplastic particles. *Scientific Reports*, 10(1), 1-12. <https://doi.org/10.1038/s41598-020-64464-9>

Ismanto, A., Hadibarata, T., Kristanti, R. A., Sugianto, D. N., Widada, S., Atmodjo, W., Satriadi, A., Anindita, M. A., Al-Mohaimeed, A. M., & Abbasi, A. M. (2023). A novel report on the occurrence of microplastics in Pekalongan River Estuary, Java Island, Indonesia. *Marine Pollution Bulletin*, 196(October), 115563. <https://doi.org/10.1016/j.marpolbul.2023.115563>

Iswanto, B., Suprapto, R., Marnis, H., & Imron, I. (2016). Performa reproduksi ikan lele Mutiara (*Clarias gariepinus*). *media akuakultur*, 11(1), 1-9. <https://doi.org/10.15578/ma.11.1.2016.1-9>

Jones, J. I., Vdovchenko, A., Cooling, D., Murphy, J. F., Arnold, A., Pretty, J. L., Spencer, K. L., Markus, A. A., Vethaak, A. D., & Resmini, M. (2020). Systematic analysis of the relative abundance of polymers occurring as microplastics in freshwaters and estuaries. *International Journal of Environmental Research and Public Health*, 17(24), 1-12. <https://doi.org/10.3390/ijerph17249304>

Kurniawan, M. A., Nugroho, S., Adnan, F., & Zulya, F. (2023). Analisis keterkaitan kelimpahan mikroplastik dengan keberadaan sampah plastik di Sungai Mahakam, Kecamatan Muara Kaman. *Jurnal Teknologi Lingkungan UNMUL*, 7(1), 20. <https://doi.org/10.30872/jtlunmul.v7i1.10822>

Li, X., Bao, L., Wei, Y., Zhao, W., Wang, F., Liu, X., Su, H., & Zhang, R. (2023). Occurrence, bioaccumulation, and risk assessment of microplastics in the aquatic environment: A review. *Water (Switzerland)*, 15(9), 1-15. <https://doi.org/10.3390/w15091768>

Liu, G., Ito, T., Kijima, Y., Yoshitake, K., Asakawa, S., Watabe, S., & Kinoshita, S. (2022). Zebrafish *Danio rerio* myotomal muscle structure and growth from a spatial transcriptomics perspective. *Genomics*, 114(5), 110477. <https://doi.org/10.1016/j.ygeno.2022.110477>

Lofty, J., Ouro, P., & Wilson, C. A. M. E. (2023). Microplastics in the riverine environment: Meta-analysis and quality criteria for developing robust field sampling procedures. *Science of the Total Environment*, 863(November 2022), 160893. <https://doi.org/10.1016/j.scitotenv.2022.160893>

Majid, D., Prabowo, A. R., Al-Kholif, M., & Sugito, S. (2018). Sintesis ferrat sebagai pendegradasi senyawa turunan benzena. *JPSE (Journal of Physical Science and Engineering)*, 3(1), 70-75. <https://doi.org/10.17977/um024v3i22018p070>

Miller, M. E., Hamann, M., & Kroon, F. J. (2020). Bioaccumulation and biomagnification of microplastics in marine organisms: A review and meta-analysis of current data. *PLoS ONE*, 15, 1-25. <https://doi.org/10.1371/journal.pone.0240792>

Moeneck, D. Y. J. A., & Toelle, N. N. (2021). GAMBARAN histologi daging ayam kampung (*Gallus domesticus*) yang diberikan ekstrak daun belimbing wuluh (*Averrhoa bilimbi*). *Jurnal Kajian Veteriner*, 9(3), 164-170. <https://doi.org/10.35508/jkv.v9i3.5596>

Nainggolan, D. H., Indarjo, A., & Suryono, C. A. (2022). Mikroplastik yang ditemukan di Perairan Karangjahe, Rembang, Jawa Tengah. *Journal of Marine Research*, 11(3), 374-382. <https://doi.org/10.14710/jmr.v11i3.35021>

Nurhadi, T., Budiyantoro, C., & Sosiasi, H. (2017).

Identifikasi mechanical properties dari bahan daur ulang polystyrene. *JMPM (Jurnal Material Dan Proses Manufaktur)*, 1(1), 36–40. <https://doi.org/10.18196/jmpm.v1i1.2758>

Nurwahyunani, A., Rakhmawati, R., & Cucianingsih, C. (2022). Kelimpahan mikroplastik pada organ pencernaan ikan Mujair (*Oreochromis mossambicus*) di Waduk Malahayu Kabupaten Brebes. *Titian Ilmu: Jurnal Ilmiah Multi Sciences*, 14(1), 18–22. <https://doi.org/10.30599/jti.v14i1.1185>

Owowenu, E. K., Nnadozie, C. F., Akamagwuna, F., Noundou, X. S., Uku, J. E., & Odume, O. N. (2023). A critical review of environmental factors influencing the transport dynamics of microplastics in riverine systems: implications for ecological studies. *Aquatic Ecology*, 57(2), 557–570. <https://doi.org/10.1007/s10452-023-10029-7>

Popi, S., Purwanto, E., Fajri, N. El, & Budijono, B. (2025). Kandungan mikroplastik pada lambung dan usus ikan Guppy (*Poecilia reticulata*) di Waduk PLTA Koto Panjang, Kabupaten Kampar, Riau, Indonesia. *Ilmu Perairan (Aquatic Science)*, 13(1), 89–95. <https://doi.org/10.31258/jipas.13.1.89-95>

Purnama, D., Johan, Y., Wilopo, M. D., Renta, P. P., Sinaga, J. M., Yosefa, J. M., & M, H. M. (2021). Analisis mikroplastik pada saluran pencernaan ikan Tongkol (*Euthynnus affinis*) hasil tangkapan nelayan di Pelabuhan Perikanan Pulau Baai Kota Bengkulu. *Jurnal Enggano*, 6(1), 110–124. <https://doi.org/10.31186/jenggano.6.1.110-124>

Roch, S., Friedrich, C., & Brinker, A. (2020). Uptake routes of microplastics in fishes: Practical and theoretical approaches to test existing theories. *Scientific Reports*, 10(1), 1–12. <https://doi.org/10.1038/s41598-020-60630-1>

Rofiq, A. A., & Sari, I. K. (2022). Analisis mikroplastik pada saluran pencernaan dan insang ikan di Brantas, Jawa Timur. *Environmental Pollution Journal*, 2(1), 263–272. <https://doi.org/10.58954/epj.v2i1.38>

Santikanuri, A. M., Haribowo, R., & Wahyuni, S. (2025). Correlation analysis of water quality and microplastic identification in the North Coast Area of Situbondo. *Jurnal Penelitian Pendidikan IPA*, 11(5), 388–397. <https://doi.org/10.29303/jppipa.v11i5.10989>

Sawalman, R., Zamani, N. P., Werorilangi, S., & Ismet, M. S. (2021). Akumulasi Mikroplastik pada spesies ikan ekonomis penting di Perairan Pulau Barrangloppo, Makassar. *Jurnal Ilmu Dan Teknologi Kelautan Tropis*, 13(2), 241–259. <https://doi.org/10.29244/jitkt.v13i2.34587>

Setyono, B. D. H., Saomadia, B. T., & Affandi, R. I. (2024). The Effect of exposure to polystyrene microplastics in feed on the growth of Tilapia (*Oreochromis niloticus*). *Jurnal Penelitian Pendidikan IPA*, 10(12), 10124–10132. <https://doi.org/10.29303/jppipa.v10i12.7924>

Smith, M., Love, D. C., Rochman, C. M., & Neff, R. A. (2018). Microplastics in seafood and the implications for human health. *Current Environmental Health Reports*, 5(3), 375–386. <https://doi.org/10.1007/s40572-018-0206-z>

Stapleton, M. J., & Hai, F. I. (2023). Microplastics as an emerging contaminant of concern to our environment: A brief overview of the sources and implications. *Bioengineered*, 14(1), 2244754. <https://doi.org/10.1080/21655979.2023.2244754>

Suwartiningsih, N., Putri, A. L. A., Wijayanti, D. E., & Prabakusuma, A. S. (2023). Deteksi Mikroplastik pada Udang Vannamei (*Littopenaeus vannamei*) dari Pasar Beringharjo Kota Yogyakarta. In *Proceeding Biology Education Conference: Biology, Science, Environmental, and Learning*, 20(1), 146–151. Retrieved from <https://jurnal.uns.ac.id/prosbi/article/view/82713>

Suwiti, N. K., Suastika, I. P., Swacita, I. B. N., & Besung, I. N. K. (2015). Histological and histomorphometry study of Bali cattle and Wagyu beef). *Jurnal Veteriner*, 16(3), 432–438. Retrieved from <https://ojs.unud.ac.id/index.php/jvet/article/view/16259>

Tang, M., Shao, T., Ma, L., Dai, H., Yong, Y., Feng, X., & Zhang, Y. (2024). Textural dynamic crisping mechanism of Crisp grass carp: Oxidative stress, apoptosis, collagen filling, and cross-linking. *Food Frontiers*, 5(2), 789–801. <https://doi.org/10.1002/fft2.354>

Utomo, E. A. T., & Muzaki, F. K. (2022). Bioakumulasi mikroplastik pada daging ikan Nila (*Oreochromis niloticus*) di Keramba Jaring Apung Ranu Grati, Pasuruan, Jawa Timur. *Jurnal Sains Dan Seni ITS*, 11(5), 26–33. <https://doi.org/10.12962/j23373520.v11i5.106895>

Wahid, A., & Joesidawati, M. I. (2024). Pencemaran mikroplastik pada tiga titik tempat pelelangan ikan di Jawa Timur. *Jurnal Ruaya*, 12(1), 85–96. <https://doi.org/10.29406/jr.v12i1>

Yang, H., Xiong, H., Mi, K., Xue, W., Wei, W., & Zhang, Y. (2020). Toxicity comparison of nano-sized and micron-sized microplastics to Goldfish *Carassius auratus* Larvae. *Journal of Hazardous Materials*, 388(January), 122058. <https://doi.org/10.1016/j.jhazmat.2020.122058>

Yona, D., Harlyan, L. I., Fuad, M. A. Z., Prananto, Y. P., Ningrum, D., & Evitantri, M. R. (2021).

Microplastic composition of *Sardinella lemuru* organs landed in Sendangbiru Port, Malang. *Journal of Fisheries and Marine Research*, 5(3), 675-684.  
<https://doi.org/10.21776/ub.jfmr.2021.005.03.20>

Yona, D., Mahendra, B. A., Fuad, M. A. Z., Sartimbul, A., & Sari, S. H. J. (2022). Kelimpahan mikroplastik pada insang dan saluran pencernaan ikan Lontok *Ophiocara porocephala* Valenciennes, 1837 (Chordata: Actinopterygii) di ekosistem mangrove Dubibir, Situbondo. *Jurnal Kelautan Tropis*, 25(1), 39-47. <https://doi.org/10.14710/jkt.v25i1.12341>

Yuranda, F., Zulkarnaini, Z., Silvia, S., & Ad'hani, C. D. (2024). Analysis of microplastics on digestive and respiration system of Horse-Eye Jack Fish (*Caranx latus*) in Coastal Bungus Bay, West Sumatra. *Jurnal Penelitian Pendidikan IPA*, 10(5), 2687-2695. <https://doi.org/10.29303/jppipa.v10i5.7099>

Zhang, L., Wu, H. X., Li, W. J., Qiao, F., Zhang, W. B., Du, Z. Y., & Zhang, M. L. (2023). Partial replacement of soybean meal by Yellow mealworm (*Tenebrio molitor*) meal influences the flesh quality of Nile tilapia (*Oreochromis niloticus*). *Animal Nutrition*, 12, 108-115.  
<https://doi.org/10.1016/j.aninu.2022.09.007>

Zhang, Z. Y., Jiang, Z. Y., Lv, H. B., Jin, J. Y., Chen, L. Q., Zhang, M. L., Du, Z. Y., & Qiao, F. (2021). Dietary aflatoxin impairs flesh quality through reducing nutritional value and changing myofiber characteristics in Yellow catfish (*Pelteobagrus fulvidraco*). *Animal Feed Science and Technology*, 274(May), 114764.  
<https://doi.org/10.1016/j.anifeedsci.2020.114764>

Zulfadhli, Z., Wijayanti, N., & Retnoaji, B. (2016). Perkembangan ovarium ikan Wader Pari (*Rasbora lateristriata* bleeker, 1854): Pendekatan histologi. *Jurnal Perikanan Tropis*, 3(1), 32-39. <https://doi.org/10.35308/jpt.v3i1.34>