



Ex-situ Conservation Efforts to the Seedlings Collected from Botanical Exploration in Mount Gede Pangrango National Park, Indonesia

Imawan Wahyu Hidayat^{1*}, Fitri Kurniawati¹, Yudi Suhendri²

¹ Research Center for Plant Conservation, Botanical Gardens and Forestry, National Research and Innovation Agency (BRIN), Bogor, Indonesia.

² Directorate of Scientific Collection Management - Cibodas Botanic Gardens, National Research and Innovation Agency (BRIN), Cipanas - Cianjur, Indonesia.

Received: July 9, 2022

Revised: April 17, 2023

Accepted: April 27, 2023

Published: April 30, 2023

Corresponding Author:

Imawan Wahyu Hidayat

imawan.wahyu.hidayat@brin.go.id

DOI: [10.29303/jppipa.v9i4.1873](https://doi.org/10.29303/jppipa.v9i4.1873)

© 2023 The Authors. This open access article is distributed under a (CC-BY License)



Abstract: A botanical exploration and collection were already conducted in March 2019 in Pasir Banteng and Pasir Pogor, Mount Gede Pangrango National Park, Indonesia, by a staff team from Cibodas Botanic Gardens (CBG). The following steps are to ensure these plants will be vigorous and survive in the nursery, which will later be ready to be planted and displayed as a garden collection. This study aimed to describe a series of treatments and management of these collections, analyze the survivorship level in the nursery, and record which seedlings were successfully planted in the garden. We explained the plant maintenance used a qualitatively-descriptively method; the survivorship level was measured based on the proportional numbers of the survived individuals; the planted seedlings in the garden were recorded from the field observation. The findings showed that in the first three months of maintenance, seedlings occurred a large amount of mortality. Next, from June 2019 to August 2020, the seedlings performed good development with no mortality reported. In September 2020, as many as 13 specimens experienced mortality. At the end of December 2020, the total crude mortality was 41 specimens, and the survivorship level was 75.15%. All the orchid specimens have been planted as the garden collections.

Keywords: Cibodas Botanic Gardens; Ex-situ Conservation; Living Plant Collection; Seedling; Survivorship.

Introduction

Human disturbances have caused severe damage to the ecosystems and human survival, including habitat loss, spreading pathogens, waste and pollution, climate changes, and many other disadvantages. Around 40% of the world's plants are threatened and face extinction risk (Antonelli et al., 2020; Mounce et al., 2017; Murphy & Romanuk, 2014). Even though various preventive measures have been adopted by nations worldwide during the last decade, the aim of halting biodiversity loss has not been achieved (Secretariat of the Convention on Biological Diversity, 2020). The plant diversity loss might lead to destruction for biological types, including humans, so more powerful maintenance of threatened

plants is urgently needed (Brummitt et al., 2015). Two key strategies for conserving plant diversity are in-situ conservation and ex-situ conservation. Ex-situ conservation is a crucial complementary purpose of in-situ conservation and may cover up the limitations of in-situ conservation (Braverman, 2014).

Ex-situ conservation means displacing living plants or seeds from their natural habitat to a man-made environment for protection and maintenance, to avoid plants from being harmed by natural causes or human disturbances (Abeli et al., 2020). Ex-situ plant collections are becoming more and more crucial for maintaining biodiversity in the face of mounting human stresses (Smith & Pence, 2017), including excessive harvesting, natural environment conversion, and climate change. Furthermore, ex-situ stocks resulting from its propaga-

How to Cite:

Hidayat, I.W., Kurniawati, F., & Suhendri, Y. (2023). Ex-situ Conservation Efforts to the Seedlings Collected from Botanical Exploration in Mount Gede Pangrango National Park, Indonesia. *Jurnal Penelitian Pendidikan IPA*, 9(4), 1873–1880. <https://doi.org/10.29303/jppipa.v9i4.1873>

tion can provide seedlings required for species reintroduction, redeployment, or supplementation (Thomas et al., 2022). The Global Strategy for Plant Conservation also trusts to enhance the value of ex-situ conservation by setting a goal of a minimum of 75% of the threatened species to preserve such as ex-situ, and 20% prepared for restoration and recovery (Sharrock et al., 2018).

Botanic gardens hold a crucial role in ex-situ conservation and a properly positioned to organize improving numbers of plant species related to conservation notice (Westwood et al., 2021). In Indonesia, one government institution obligated to establish and conduct ex-situ conservation efforts is the botanical gardens. Based on Presidential Regulation Number 93 of 2011, botanical gardens have some duties to carry out ex-situ plant conservation efforts and manage the site that contains plant collection resulting from these efforts, and the data must be well-documented.

Cibodas Botanic Gardens (CBG) is an Indonesian government institution as one of the botanic gardens under the National Research and Innovation Agency (*Badan Riset dan Inovasi Nasional - BRIN*) of the Republic of Indonesia. CBG is concerned with conducting ex-situ conservation of wet-tropic montane plants, particularly from western Indonesia. Tropical montane plants are urgent to be conserved due to the rate of degradation in these regions also tends to increase (Boehmer, 2011). Worldwide climate changes are also a potential warning in the conservation efforts regarding montane plants (Rathore et al., 2019; Thang et al., 2020). Therefore, CBG needs to conduct simultaneous efforts concerning ex-situ montane plant conservation.

In March 2019, a CBG team conducted a botanical exploration and collection in Pasir Banteng and Pasir Pogor, Mount Gede Pangrango National Park, Indonesia. Within those efforts, hundreds of specimens from 36 plant families were successfully collected and transferred to the nursery of CBG (Hidayat & Kurniawati, 2021). The next critical steps are to ensure the survivorship of these collections to keep thriving and vigor, so the latter are ready to be planted in the garden and become plant collections.

Ensuring the seedlings keep healthy and flourishing needs extra attention and maintenance. The level of seedling mortality may reach 90% or even more during their early development. Therefore, this period is a critical life phase of the tree seedlings (Conlisk et al., 2017). So, this study aimed to describe a series of treatments of these collections in the nursery, analyze the survivorship of the seedlings, and record the seedlings that were successfully planted as the garden living plant collection. The findings will hopefully increase the understanding of the value of ex-situ plant conservation efforts, benefiting future research, educational endeavors, and public awareness.

Method

A botanical exploration and collection were already conducted in Pasir Banteng and Pasir Pogor, Mount Gede Pangrango National Park, Indonesia, in March 2019. The activities successfully collected as many as 165 specimens (53 species from 36 families) in the form of seedlings (Hidayat & Kurniawati, 2021). Then, all the seedlings collected were delivered to the nursery unit of CBG to receive acclimatization and maintenance treatments.

The process and survivorship observation were conducted from the end of March 2019 to December 2020. The intervals were considered to be trustworthy and capable of enhancing seedling survival. After these periods, the maintenance in the nursery keeps on going until the seedlings are ready to be planted in the garden. However, it will not be further explained. Then, the recording of which seedlings were planted as a garden collection was observed until December 2020.

Preparation Stage

The nursery unit of CBG has been prepared with a greenhouse to accommodate the arrived seedlings from the field. To avoid direct exposure to the sunlight, keep the moisture, and reduce the transpiration, then at the top of the inner room was installed a paranet shade with a density of 65% (Irawan et al., 2020).

Each seedling will be planted in a polybag of diameter (\emptyset) of 15 cm in size. A growth medium of the tree seedlings and terrestrial orchids was a mix of forest compost and paddy husks with a ratio of 1:1 (Irawan et al., 2020; Prasetyonohadi & Kuswanto, 2016). Epiphyte orchids, such as *Bulbophyllum* spp., were grown in the medium of chopped ferns. The different mediums used to adapt the seedlings to be compatible with their natural habitat in the wild. The mediums were also given an insecticide (*Furadan*®) at a dose of 7.5 to 10 gr. each polybag to prevent interference from worms, termites, and other insects.

Maintenance Stage

Soon the seedlings arrived at the nursery, each plant displaced to the polybag added with medium according to their types (tree, terrestrial orchid, or epiphyte orchid). Only *Bulbophyllum* sp. was grown in the medium of chopped ferns, and the rest were planted in a mix of forest compost and paddy husks medium. Each specimen was already moved to a polybag, then equipped with a plant identity (i.e., species name, plant family name, the collector's name code, date of collection, and elevation).

All polybags were well-arranged on the greenhouse floor which allows the staff's movement during the treatment. Then, it was conducted by watering once to twice a day, depending on the room's temperature and

humidity. Weed and pest control once a week, growth hormone supplies once in two weeks, and fertilizing once a month. Growth hormone was added to restore the plant condition and stimulate shoots and root development (Irawan et al., 2020). These serial processes are conducted to the seedlings at the nursery unit until they are ready and vigor planted in the garden. However, in this study, the maintenance observation was only conducted from the end of March 2019 to December 2020.

To calculate the survivorship level of the seedlings during these periods, then be analyzed some parameters, namely: crude mortality rate (d), which is the ratio of the number of deaths during a specific period (in this study was twenty-one months) to the number of the previous population; proportional mortality rate (q), which is the proportion of death individuals to the number of the early population; and, survivorship level (l), which is the proportion of the survived individuals to the number of the early population at the end of the observation period (Alvarez-Aquino & Williams-Linera, 2012). The survivorship observation was conducted once a week and accumulated monthly. Next, all the parameters are presented in tables and graphics.

Garden Plant Collection Record

After the seedlings were maintained and morphologically thrive and vigorous, they were ready to be planted in the garden. The planting location of new plants is usually based on the closeness of kinship or taxon with other plants. It also considers the adequacy of sunlight, soil type, topography soil, and water availability. Then, after three months in the garden field, if the plant performs good development, it is registered as a new plant seedling collection. Next, this new data is recorded in the garden collections database.

Result and Discussion

In the early three months of maintenance, it occurred a significant loss of the seedlings. As many as 28 specimens were dead that 13 specimens in May and 15 specimens in June 2019 (Table 1). Even *Lindera polyantha* (Lauraceae) and *Lindsaea cultrata* (Lindsaeaceae) were unable to survive in their first three months. The species of *L. polyantha* has a high ability in water absorption (Lailati, 2021). Despite its tolerance to wood decay being moderately resistant Djarwanto et al. (2019), the CBG's microclimates which are characterized by high to very high humidity (> 85%), caused the failure of the early development stage of these seedlings. For *L. cultrata*, it was assumed that the growth medium and the microclimates were unsuitable likely their natural habitat in the deep forest (Abas, 2017).

It was 17 species (from 15 families) that experienced mortality in their population (d). In this first critical

period of plant seedling development, a plant frequently has some difficulties to well-grow or even often dies (Conlisk et al., 2017). At this early stage of the seedling development, the survivorship level (l) reached 83.03% accumulatively, and the proportional mortality rate (q) reached 16.97%, or there occurred ~17 specimens' mortality in every 100 populations.




Figure 1. Survivorship Curve of the Total Seedlings in their First Three Months of Maintenance

Figure 1 also describes the survivorship condition of the seedling in these first three months of maintenance. The coefficient was negative, and the value of R^2 was 96.39%. It means that the number of mortalities is intensely significant in affecting the survivorship level of the seedling population. At this critical period, seedlings are highly sensitive to drought and desiccation. Nursery conditions also roles a significant part in emerging resilience (Palma & Laurance, 2015). For example, the facility of the watering system and microclimate control, especially temperature and air humidity, should fulfill the seedling's necessity to grow.

The other species occurred mortality even though not the entire specimen died, at least leaving one or two specimens to live. Some of these reasons were a new condition at the CBG's nursery is a shocking environment for these new seedlings. A recent circumstance, such as microclimates and a new growth medium, are unable to support its early development or vice versa, and these new seedlings failed to adapt to these new conditions. Microclimates supported by other biophysical factors have a linear influence on plant growth and sustainability (Maclean et al., 2015).

The growth medium in composition between forest compost and paddy husk with a ratio of 1:1 still needs further research. An accurate and suitable growth medium may be different among plant seedlings because each plant requires a specific growth medium and composition, such as soil pH (Offord et al., 2014).

The different quality of each seedling from its natural habitat is also assumed to be a driving factor of the failure. For certain species, seedlings of stressed resistant grandparents or parents generated a faster regeneration, more developed root system, and growing

faster than seedlings from the same genetic lineages whose grandparents or parents did not receive stress conditions (Herman et al., 2012; Palma & Laurance, 2015).

Table 1. The Seedlings Experienced Mortality in Their First Three Months of Maintenance

Species name	Family	Early population (n_0)	Survived population (n_i)			Crude mortality (d_i)
			2019			
			April	May	June	
<i>Acronychia trifoliolata</i> Zoll. & Moritzi	Rutaceae	3	3	2	1	2
<i>Begonia muricata</i> Blume	Begoniaceae	5	5	4	3	2
<i>Cryptocarya</i> sp.	Lauraceae	2	2	1	1	1
<i>Cryptostylis arachnites</i> (Blume) Hassk.	Orchidaceae	4	4	3	3	1
<i>Daphniphyllum glaucescens</i> Blume	Daphniphyllaceae	3	3	3	1	2
<i>Eurya acuminata</i> DC.	Pentaphylacaceae	4	4	4	2	2
<i>Fagraea elliptica</i> Roxb.	Gentianaceae	3	3	2	2	1
<i>Lindera polyantha</i> Boerl.	Lauraceae	1	1	1	0	1
<i>Lindsaea cultrata</i> (Willd.) Sw.	Lindsaeaceae	5	5	3	0	5
<i>Polyalthia subcordata</i> (Blume) Blume	Annonaceae	5	5	4	3	2
<i>Schima wallichii</i> (DC.) Korth.	Theaceae	5	5	5	4	1
<i>Strobilanthes paniculata</i> (Nees) Miq.	Acanthaceae	6	6	5	4	2
<i>Turpinia montana</i> (Blume) Kurz	Staphyleaceae	4	4	3	3	1
<i>Urophyllum</i> sp.	Rubiaceae	4	4	3	3	1
<i>Strobilanthes involucrata</i> Blume	Acanthaceae	5	5	5	3	2
<i>Strobocalyx arborea</i> (Buch.-Ham.) Sch.Bip.	Compositae	2	2	1	1	1
<i>Viburnum sambucinum</i> Reinw. ex Blume	Adoxaceae	2	2	1	1	1

Furthermore, from June 2019 to August 2020, the seedlings started to perform good adaptation and development in the new environment. There was no death of the seedlings reported in that period. It indicated that the treatments and maintenance efforts performed by CBG's staff were already suited to the plant growth needs. Moreover, in July 2020, all the orchid species have been displaced to the orchid house and planted as garden collections (Table 2).

Table 2. Orchid Species Planted as Garden Collections

Orchid species	Genus	Number of individuals
<i>Apostasia wallichii</i> R.Br.	Apostasia	3
<i>Bulbophyllum</i> sp.	Bulbophyllum	5
<i>Calanthe</i> sp.	Calanthe	5
<i>Cryptostylis arachnites</i> (Blume) Hassk.	Cryptostylis	2
<i>Cymbidium lancifolium</i> Hook.	Cymbidium	4
<i>Plocoglottis javanica</i> Blume	Plocoglottis	5

Environment degradations have several impacts on the species variety, including orchids. Due to overharvesting, orchids are one of the most endangered flowering plants. However, orchid species are going extinct in the wild caused of their lost habitat. Humans' intrusions and activities, such as agricultural practices, urbanization, extending settlements, deforestation, and illegal harvesting for economic purposes, are threatened

the existence of orchid species in their natural habitat (Phillips et al., 2020; Seaton et al., 2013). Therefore, orchids are crucial to be preserved and as one of the main targeted plants in every botanical exploration and conservation activity conducted by CBG.

Next, in September 2020, several mortalities occurred in the seedlings. As many as 13 specimens from nine species occurred mortality in their population (Table 3). *Calophyllum* sp., *Rubus alceifolius*, and *Strobocalyx arborea* experienced complete death and left no surviving seedling. It is a bad fortune, even though this species has potential uses as an ornamental and medicinal plant (Slik, 2022; Valkute et al., 2018).

The most common matter assumed for these failures was stem and root decay. Highly humidity surrounding the nursery has implications for humidity conditions in the greenhouse. During the seedling development, it should be a good humidity circulation in the greenhouse. Where moisture losses from ventilation and the stability between plant transpiration and soil evapotranspiration (Shamshiri et al., 2018). Microclimate conditions around the greenhouse, such as low temperatures combined with high humidity at night and in the morning, and non-optimal air circulation in the greenhouse, are assumed to result in water vapor excess that causes the decayed of the seedlings.

Table 3. The Seedlings Experienced Mortality Between August to September 2020

Species name	Family	Survived population (n_i)		Crude mortality (d_i)
		August	September	
<i>Ardisia villosa</i> Roxb.	Primulaceae	5	4	1
<i>Calophyllum</i> sp.	Clusiaceae	2	0	2
<i>Eriosolena composita</i> (L.f.) Tiegh.	Thymelaeaceae	2	1	1
<i>Medinilla speciosa</i> Blume	Melastomataceae	3	1	2
<i>Strobilanthes involucreta</i> Blume	Acanthaceae	3	2	1
<i>Myrsine avenis</i> (Blume) A.DC.	Primulaceae	2	1	1
<i>Rubus alceifolius</i> Poir.	Rosaceae	2	0	2
<i>Tacca chantrieri</i> André	Dioscoreaceae	3	1	2
<i>Strobocalyx arborea</i> (Buch.-Ham.) Sch.Bip.	Compositae	1	0	1

Furthermore, from September to the end of December 2020, the number of seedlings remain stable, with no mortality experience. Until this period, no additional seedlings have been planted in the garden for collection. These seedlings remain maintained in the nursery, awaiting the upcoming planting schedule.

Overall, during the maintenance periods (from April 2019 to December 2020), the seedling population experienced mortality or crude mortality (d) of as many as 41 specimens. Survivorship level (l) reached 75.15%, where R^2 was 69% (Figure 2). It means that the population decreases are significantly influenced by the amount of mortality. The proportional mortality rate (q) was 24.85%, or there occurred ~25 specimens' mortality in every 100 populations. However, this level was considered a moderate success in managing the seedlings survivorship. Some important notes that need attention are how the best manage or modify the microclimate in the greenhouse so that it remains to fulfill the requirements for optimal seedling growth.




Figure 2. Survivorship Curve of The Total Seedlings After Twenty-One Months of Maintenance

Based on the type of the curve, the survivorship characteristic of this study included the type of III. This type describes how the number of mortalities at the early stage of development tends to be high, but then the life expectancy gradually improves for the individuals who survived at their first stage (Molles & Sher, 2019). This

description corresponds with this study. In the first three months, plant seedlings suffered high mortality, but then in the following periods, the mortality level decreased, even to zero. It was only experienced another mortality with a lower amount than the previous. At this stage, the plants tried to adapt to a new environment after being displaced from their natural habitat. The differences in the microclimate and other biophysical conditions influence the survival rate of the seedlings.




Figure 3. Survivorship Proportion Between the Number of Initial and Survived Seedlings Based on Their Families

This study also showed that specimens from families of Clusiaceae, Compositae, Lindsaeaceae, and Rosaceae failed to survive during the maintenance pe-

riod (Figure 3). They contributed as much as 26.83% of the total mortality. The remaining families were left with at least a specimen to be maintained in the nursery, such as Adoxaceae, Chloranthaceae, Daphniphyllaceae, Dioscoreaceae, Lauraceae, Melastomataceae, Oleaceae, and Thymelaeaceae. As many as 15 families that showed 100% successfully survived during these periods, such as Araceae, Chloranthaceae, Dichapetalaceae, Elaeocarpaceae, Escalloniaceae, etc.

The optimization efforts to maintain the remaining surviving seedlings still be continued. Watering, fertilizing, providing the supplement, and weed and pest control are routinely performed, either daily or weekly. Hopefully, with these routine maintenances along with plant maturation, the survival rate might be increased or at least be maintained. The seedlings will keep placed at the nursery until later planting activities are ready to be established and the seedlings confirmed planted in the garden.

Based on the above description, ex-situ plant conservation conducted on those seedlings gives a crucial role in the plant stock availability of CBG to enrich its collections. Plant enrichment is not only significant to the existence of CBG but also enhances the value of plant conservation. When those seedlings are eventually planted in the garden, they may trigger further research and educational purposes that are beneficiaries to the community. These efforts are also a form of plant species preservation and rescue when natural habitats continuedly experiencing degradation and destruction problems.

Conclusion

At the early stage of development, the seedlings experienced significant losses. But then, the maintenance efforts gradually succeeded in reducing the losses and keeping the survivorship relatively stable. The nursery staffs of CBG were capable of maintaining the survivorship of three-quarters of the total seedlings. All the orchid seedlings were planted as garden collections and displaced to the orchid's house. However, this study pointed out that the greenhouse facilities and the knowledge of the proper growing media need to be developed immediately to improve the survivorship level of the maintained seedlings. The survived seedlings were a crucial plant stock to enrich CBG's plant collections and enhance the value of ex-situ plant conservation, particularly for the tropical montane plants.

Acknowledgements

The plant seedlings management and field observation were partly funded by *DIPA Tematik* of the Indonesian Institute of Sciences (LIPI) in the 2019 fiscal year. We are

grateful to Muslim, Rustandi A., Nudin, Agus Supratman, Emus, and all staff of CBG's nursery unit who conducted maintenance of the seedlings.

References

- Abas, C. G. S. (2017). Ferns and Lycophytes as indicators of forest environment of Kampo Uno, Katipunan, Davao-Arakan Valley, North Cotabato. *International Journal of Engineering, Science and Mathematics*, 6(8), 10-16. Retrieved from <https://www.indianjournals.com/ijor.aspx?target=ijor:ijesm&volume=6&issue=8&article=002>
- Abeli, T., Dalrymple, S., Godefroid, S., Mondoni, A., Müller, J. V., Rossi, G., & Orsenigo, S. (2020). Ex situ collections and their potential for the restoration of extinct plants. *Conservation Biology*, 34(2), 303-313. <https://doi.org/10.1111/cobi.13391>
- Alvarez-Aquino, C., & Williams-Linera, G. (2012). Seedling survival and growth of tree species: Site condition and seasonality in tropical dry forest restoration. *Botanical Sciences*, 90(3), 341-351. <https://doi.org/10.17129/botsci.395>
- Antonelli, A., Fry, C., Smith, R. J., Simmonds, M. S. J., Kersey, P. J., Pritchard, H. W., ... Zhang, B. G. (2020). *State of the World's Plants and Fungi*. Doctoral dissertation, Royal Botanic Gardens (Kew); Sfumato Foundation. <https://doi.org/10.34885/172>
- Boehmer, H. J. (2011). Vulnerability of tropical montane rain forest ecosystems due to climate change. In *Coping with Global Environmental Change, Disasters and Security: Threats, Challenges, Vulnerabilities and Risk*, 5, 789-802. https://doi.org/10.1007/978-3-642-17776-7_46
- Braverman, I. (2014). Conservation without nature: The trouble with in situ versus ex situ conservation. *Geoforum*, 51, 47-57. <https://doi.org/10.1016/j.geoforum.2013.09.018>
- Brummitt, N. A., Bachman, S. P., Griffiths-Lee, J., Lutz, M., Moat, J. F., Farjon, A., ... Lughadha, E. M. N. (2015). Green plants in the red: A baseline global assessment for the IUCN Sampled Red List Index for Plants. *PLoS ONE*, 10(8), 1-22. <https://doi.org/10.1371/journal.pone.0135152>
- Conlisk, E., Castanha, C., Germino, M. J., Veblen, T. T., Smith, J. M., & Kueppers, L. M. (2017). Declines in low-elevation subalpine tree populations outpace growth in high-elevation populations with warming. *Journal of Ecology*, 105(5), 1347-1357. <https://doi.org/10.1111/1365-2745.12750>
- Djarwanto, Suprapti, S., Abdurrachman, & Arsyad, W. M. (2019). Fungal decay resistance of 16 tropical wood species embedded with metal screws. *Journal of Tropical Forest Science*, 31(4), 443-451. <https://doi.org/10.26525/jtfs2019.31.4.443>

- Herman, J. J., Sultan, S. E., Horgan-Kobelski, T., & Riggs, C. (2012). Adaptive transgenerational plasticity in an annual plant: Grandparental and parental drought stress enhance performance of seedlings in dry soil. *Integrative and Comparative Biology*, 52(1), 77-88. <https://doi.org/10.1093/icb/ics041>
- Hidayat, I. W., & Kurniawati, F. (2021). Botanical exploration and conservation in Pasir Banteng and Pasir Pogor, Gunung Gede Pangrango National Park. *Jurnal Sylva Lestari*, 9(3), 488-502. <https://doi.org/10.23960/jsl.v9i3.540>
- Irawan, U., Arbainsyah, Ramlan, A., Putranto, H., & Afifudin, S. (2020). *Manual Pembuatan Persemaian Dan Pembibitan Tanaman Hutan*. Bogor: Operasi Wallacea Terpadu. [Indonesian]. Retrieved from https://elti.yale.edu/sites/default/files/rsource_files/buku_manual_persemaian_dan_pembibitan_tanaman_hutan.pdf
- Lailati, M. (2021). Kajian beberapa tanaman dataran tinggi koleksi Kebun Raya Cibodas dalam kemampuan penyerapan dan konservasi air. *Prosiding Seminar Nasional Biologi (SEMABIO) 6*, 6(1), 79-87. Retrieved from <https://conferences.uinsgd.ac.id/index.php/gdcs/article/view/494>
- Maclean, I. M. D., Hopkins, J. J., Bennie, J., Lawson, C. R., & Wilson, R. J. (2015). Microclimates buffer the responses of plant communities to climate change. *Global Ecology and Biogeography*, 24(11), 1340-1350. <https://doi.org/10.1111/geb.12359>
- Molles, M. C., & Sher, A. A. (2019). *Ecology: Concepts and Applications* (8th ed.). New York: McGraw-Hill Education.
- Mounce, R., Smith, P., & Brockington, S. (2017). Ex situ conservation of plant diversity in the world's botanic gardens. *Nature Plants*, 3(10), 795-802. <https://doi.org/10.1038/s41477-017-0019-3>
- Murphy, G. E. P., & Romanuk, T. N. (2014). A meta-analysis of declines in local species richness from human disturbances. *Ecology and Evolution*, 4(1), 91-103. <https://doi.org/10.1002/ece3.909>
- Offord, C. A., Meagher, P. F., & Zimmer, H. C. (2014). Growing up or growing out? How soil pH and light affect seedling growth of a relictual rainforest tree. *AoB PLANTS*, 6(1). <https://doi.org/10.1093/aobpla/plu011>
- Palma, A. C., & Laurance, S. G. W. (2015). A review of the use of direct seeding and seedling plantings in restoration: What do we know and where should we go? *Applied Vegetation Science*, 18(4), 561-568. <https://doi.org/10.1111/avsc.12173>
- Phillips, R. D., Reiter, N., & Peakall, R. (2020). Orchid conservation: From theory to practice. *Annals of Botany*, 126(3), 345-362. <https://doi.org/10.1093/aob/mcaa093>
- Prasetyonohadi, D., & Kuswanto, H. (2016). *Pedoman Persemaian: Sebuah Panduan Menuju Bibit yang Sehat*. Jakarta: Kementerian Kehutanan dan Lingkungan Hidup, (Indonesian).
- Rathore, P., Roy, A., & Karnatak, H. (2019). Modelling the vulnerability of *Taxus wallichiana* to climate change scenarios in South East Asia. *Ecological Indicators*, 102, 199-207. <https://doi.org/10.1016/j.ecolind.2019.02.020>
- Seaton, P., Kendon, J. P., Pritchard, H. W., Puspitaningtyas, D. M., & Marks, T. R. (2013). Orchid conservation: The next ten years. *Lankesteriana*, 13(1), 93-101. <https://doi.org/10.15517/lank.v0i0.11545>
- Secretariat of the Convention on Biological Diversity. (2020). *Global Biodiversity Outlook 5*. Montreal. Retrieved from <https://www.cbd.int/gbo/gbo5/publication/gbo-5-en.pdf>
- Shamshiri, R. R., Jones, J. W., Thorp, K. R., Ahmad, D., Man, H. C., & Taheri, S. (2018). Review of optimum temperature, humidity, and vapour pressure deficit for microclimate evaluation and control in greenhouse cultivation of tomato: A review. *International Agrophysics*, 32(2), 287-302. <https://doi.org/10.1515/intag-2017-0005>
- Sharrock, S., Hoft, R., & Dias, B. F. S. (2018). An overview of recent progress in the implementation of the Global Strategy for Plant Conservation-a global perspective. *Rodriguésia*, 69(4), 1489-1511. <https://doi.org/10.1590/2175-7860201869401>
- Slik, J. W. F. (2022). *Family Classification*. Plants of Southeast Asia. Retrieved from <https://www.asianplant.net>
- Smith, P., & Pence, V. (2017). The Role of Botanic Gardens in Ex situ Conservation. In *Plant Conservation Science and Practice: The Role of Botanic Gardens*, 102-133. Retrieved from [http://refhub.elsevier.com/S0006-3207\(21\)00462-6/1f202111240448529687](http://refhub.elsevier.com/S0006-3207(21)00462-6/1f202111240448529687)
- Thang, T. H., Thu, A. M., & Chen, J. (2020). Tree species of tropical and temperate lineages in a tropical Asian montane forest show different range dynamics in response to climate change. *Global Ecology and Conservation*, 22, e00973. <https://doi.org/10.1016/j.gecco.2020.e00973>
- Thomas, G., Sucher, R., Wyatt, A., & Jiménez, I. (2022). Ex situ species conservation: Predicting plant survival in botanic gardens based on climatic provenance. *Biological Conservation*, 265(1), 109410. <https://doi.org/10.1016/j.biocon.2021.109410>
- Valkute, T. R., Aratikatla, E. K., Gupta, N. A., Ganga, S., Santra, M. K., & Bhattacharya, A. K. (2018). Synthesis and anticancer studies of Michael adducts and Heck arylation products of sesquiterpene lactones, zaluzanin D and zaluzanin C from *Vernonia arborea*. *RSC Advances*, 8(67),

38289-38304. <https://doi.org/10.1039/c8ra06238b>
Westwood, M., Cavender, N., Meyer, A., & Smith, P.
(2021). Botanic garden solutions to the plant
extinction crisis. *Plants People Planet*, 3(1), 22-32.
<https://doi.org/10.1002/ppp3.10134>